

Trends in Utilizing CT Scan in the Emergency Department: Correlation with Significant Imaging Findings and Subsequent Hospital Admissions. A Retrospective Study

Mohammed Wazzan¹, Ahmed Abduljabbar¹, Rani Ahmed¹, Majed Alkhatib², Khaled Bahubaishi³, Hassan Alshehri⁴, Mohammad Bukhary⁵, Amr Mansouri¹, Ahmad H. Bakhribah⁶, Imad M. Khojah⁶, Mohammed Sabri¹, Mohammed Addas⁷

¹Department of Radiology, Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia

²Department of Radiology, International Medical Center, Jeddah, Saudi Arabia

³Teleradiology Section, Diagnostic Elite, Jeddah, Saudi Arabia

⁴Department of Radiology, King Fahad Armed Forces Hospital, Jeddah, Saudi Arabia

⁵Anesthesia Department, Alhada Armed Forces Hospital, Taif, Saudi Arabia

⁶Department of Emergency Medicine, Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia

⁷Department of Radiology, King Abdullah Medical Complex, Jeddah, Saudi Arabia

Correspondence

Mohammed Wazzan

Department of Radiology, Faculty of Medicine,
King Abdulaziz University
P.O. Box 80205, Jeddah 21589
Kingdom of Saudi Arabia
e-M: mawazzan@kau.edu.sa

Submission: 29 Aug. 2023

Accepted: 08 Dec. 2024

Citation

Wazzan M, Abduljabbar A, Ahmed R, Alkhatib M, Bahubaishi K, Alshehri H, Bukhary M, Mansouri A, Bakhribah AH, Khojah IM, Sabri M, and Addas M. Trends in utilizing CT scan in the emergency department: correlation with significant imaging findings and subsequent hospital admissions. A retrospective study JKAU Med Sci 2024; 31(1): 1–8. DOI: 10.4197/Med.31–1.1.

Copyright: ©The Author(s), YEAR. Publisher. The Journal of King Abdulaziz University - Medical Sciences is an Official Publication of "King Abdulaziz University". It is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Objective: To assess the association between boarding time in the emergency department (ED) and increased patient mortality at the King Abdul-Aziz University Hospital in Jeddah, Saudi Arabia.

Methods: A retrospective study was conducted on 28,066 patients admitted through the ED and discharged from January to December 2019 at the King Abdul-Aziz University Hospital in Jeddah, Saudi Arabia. Data on patients' demographics, age, nationality, arrival method, eligibility, triage level, and outcome (alive or dead) was collected from medical records.

Results: This study found a strong link between patient mortality and boarding time, with patients who died experiencing significantly longer boarding times than those who survived. The average boarding time for those who died was more than 12 hours, compared to only 2.2% of those who did not. Additionally, the death rate was higher among non-eligible and male patients. Patients with injuries or orthopedic diseases had a notably increased likelihood of boarding for more than 12 hours. Both long-term boarding and non-eligibility for management were risk factors for death among the patients studied.

Conclusion: This paper highlights the impact of protracted boarding times on patient outcomes and emphasizes the urgent need to take immediate steps to decrease ED boarding.

Keywords

Association, Boarding, Emergency, Mortality

INTRODUCTION

Emergency department (ED) overcrowding is a significant challenge for hospitals and is considered a hospital-wide problem^[1]. It has been described as “the most serious issue confronting EDs in the developed world^[2]. It is associated with poor quality of care and unfavorable events following ED assessment^[3,4], leading to delays in treatment and high complication rates, which have increased the mortality rate^[5-12]. Although unnecessary ED visits are thought to be the cause of ED overcrowding, ED boarding seems to be the leading cause, characterized by patient waiting time in the ED after primary assessment and initial care^[7, 13,14]. ED boarding can have serious implications that act as major barriers to advanced medical treatment and can lead to delays in time-sensitive procedures^[14,15]. For example, the outcome in sepsis and septic shock patients can demonstrate a substantial change if seen in early target therapy^[16,17]. Studies conducted in France, Greece, and Canada found that an increased length of stay and boarding in the ED is associated with increased mortality and comorbidity^[17,18,19]. However, a large retrospective study showed prolonged stays in the emergency department may increase the risk of mobility problems for patients but not necessarily increase motility^[20].

In the Kingdom of Saudi Arabia (KSA), ED non-emergency cases are considered the leading cause of ED overcrowding, adversely affecting healthcare providers, patients, and the country's economy^[21]. In KSA culture, patients prefer to stay in the ED for better care. A 2018 study was conducted to identify the preferences of patients and their attendants regarding staying in the ED or being transferred to inpatient units. The results indicated that 59% of patients preferred staying in the ED due to the facilities and prompt attention provided by the staff. The study emphasized the importance of a timely transfer to inpatient units to reduce medical mistakes and complications, especially in overcrowded EDs. However, until now, there has not been enough data to support the idea that there is a clear relationship between a long stay in the ED and mortality from local hospitals in KSA.

Given the controversy surrounding the issue and the limited number of studies conducted in KSA, this retrospective study aims to examine the association between boarding time in the ED and increased patient mortality at King Abdul-Aziz University Hospital in Jeddah, Saudi Arabia, in 2020–2021.

SUBJECTS AND METHODS

Study design and time frame: This study was a retrospective study conducted from 2020 to 2021.

STUDY SETTING

The study was conducted at the Emergency Department (ED) of the King Abdul-Aziz University Hospital in Jeddah, Saudi Arabia. This tertiary care center serves the middle- and low-income population in Jeddah.

STUDY POPULATION

All patients admitted to the hospital through the ED and discharged from January to December 2019 were included, leaving a sample size of 28,066 patients. Patients with missing data on the system, OB patients, and pediatrics were excluded.

TIMELINE

In this study, we will consider a stay in the ED for more than 12 hours to be a prolonged ED stay.

DATA COLLECTION

Data was collected from patients' medical records, including demographics, age, nationality, arrival method, eligibility, outcome, triage level, and whether they died or were still alive.

STATISTICAL ANALYSIS

Data was coded, tabulated, and analyzed using SPSS version 20 (Armonk, NY: IBM Corp.). Qualitative data was expressed as numbers and percentages, and the Chi-squared test (χ^2) was used to test the relationship between variables. Quantitative data was expressed as mean and standard deviation (Mean \pm SD), and the Mann-Whitney test was used for non-parametric variables. Binary logistic regression was used to assess the independent predictors of death in the ED. A p-value of < 0.05 was considered statistically significant.

RESULTS

The total number of patients enrolled in the study was 28,066 patients. Table 1 shows that the age of male participants was 34.17 ± 20.76 years, with 50.4% male and 78.7% Saudi nationals. 52.8% of patients had a triage level of priority 4 (less urgent), 99.2% arrived by

Table 1. Distribution of the studied patients according to their characters, arrival method, eligibility, outcome, triage level, and their fate

Variable	No (%)
Age (mean SD)	34.17±20.76
Boarding time (mean SD)	4.35 ± 3.79
Gender	
Female	13,919 (49.6)
Male	14,147 (50.4)
Nationality	
Saudi	22,095 (78.7)
Non - Saudi	5,971(21.3)
Triage level	
Priority 1. Resuscitation	327 (1.2)
Priority 2. Emergent	4,008 (14.3)
Priority 3. Urgent	8,039 (28.6)
Priority 4. Less Urgent	1,480 (52.8)
Priority 5. Non-Urgent	887 (3.2)
Arrival method	
Ambulance	229 (0.8)
Other (Arrival)	27,837 (99.2)
Eligible	
No	25,543 (91.0)
Yes	2,523 (9.0)
Arrival timing	
am	16,302 (58.1)
pm	11,764 (41.9)

a method other than ambulance, 91% were eligible for management in the ED, and 76.9% were discharged. For 77.1% of patients, the main responsible physician (MEP) was an emergency physician. The average boarding time was recorded as 4.35 ± 3.79 hours. Of all patient arrivals, 58.1% occurred during the day shift (7 AM to 7 PM), while 41.9% occurred during the night shift (7 PM to 7 AM). The total mortality rate was found to be 1.5% for all arrivals. The most common comorbidities among patients were heart disorders (16.3%), followed by pulmonary disorders (12.8%), and musculoskeletal disorders (12%).

Figure 1 shows that half of the participants (50.8%) had a boarding time of less than two hours, while 2.9% had a boarding time of more than 12 hours.

Table 2 shows that patients who died had a significantly higher boarding time (11.23 ± 5.56 hours) compared to those who did not die (4.24 ± 3.66 hours), with a p-value of < 0.001. 46.2% of patients who died

Boarding time (mean ± SD)	4.35 ± 3.79
Fate	
No death	27,655 (99.5)
Death	411 (1.5)
Comorbidities	
Heart disorders	4,582 (16.3)
Hypertension	3,075 (11.0)
Diabetes mellitus	2,902 (10.3)
GIT disorders	2,885 (10.3)
Pulmonary disorders	3,590 (12.8)
Neurologic disorders	2,145 (7.6)
Musculoskeletal disorders	3,376 (12.0)
Skin disorders	969 (3.5)
ENT disorders	777 (2.8)
Injury and orthopedic disorders	637 (2.3)
Urinary disorders	577 (2.1)
Thyroid disorders	653 (2.3)
Blood disorders	587 (2.1)
Fluid and electrolyte disorders	56 (0.9)
Deficiency anemias	414 (1.5)
Psychological disorders	631 (2.2)

had a boarding time of more than 12 hours, compared to only 2.2% for those who did not die (Figure 2). Furthermore, there was a significant relationship between patients' deaths and their eligibility status, with a p-value of 0.006. However, no significant relationship was found between patients' deaths and their age, nationality, gender, arrival method, arrival timing, outcome, MRP, triage level, or comorbidities with a p-value of > 0.05.

Table 3 shows that male patients had a significantly longer boarding time of more than 12 hours compared to female patients. A non-significant difference was found between boarding time and patients' age, nationality, arrival method, eligibility, outcome, MRP, arrival timing, or triage level ($p = > 0.05$). Patients who had injuries or orthopedic disorders had a significantly higher percentage of those who had boarding time of more than 12 hours (5.2%), followed by those who had neurological disorders (3.8%) and those who had pulmonary disorders (3.5%) ($p = < 0.05$).

Table 4 shows that binary logistic regression analysis found long boarding time and non-eligibility for management as independent predictors of death among studied patients.




Figure 1. Distribution of the studied patients according to their boarding time.

Table 2. Relationship between patients' death and their characters, arrival method, eligibility, outcome, triage level, and comorbidities


Variable	Death		Test	p-value
	Death	No Death		
Age	34.16 ± 20.74	35.11 ± 22.22	0.53*	0.59
Boarding time	11.23 ± 5.56	4.24 ± 3.66	24.4*	< 0.001
Gender				
Female	213 (1.5)	13706 (98.5)	0.83**	0.36
Male	198 (1.4)	13949 (98.6)		
Triage level				
Priority 1 - Resuscitation	2 (0.6)	325 (99.4)	8.46**	0.07
Priority 2 - Emergent	75 (1.9)	3933 (98.1)		
Priority 3 - Urgent	102 (1.3)	7937 (98.7)		
Priority 4 - Less Urgent	218 (1.5)	14587 (98.5)		
Priority 5 - Non-Urgent	14 (1.6)	873 (98.4)		
Nationality				
Saudi	314 (1.4)	21781 (98.6)	1.34**	0.24
Non - Saudi	97 (1.6)	5874 (98.4)		
Arrival method				
Ambulance	3 (1.3)	226 (98.7)	0.03**	0.84
Other	408 (1.5)	27429 (98.5)		
Eligibility				
No	52 (2.1)	25184 (98.6)	6.93**	0.006
Yes	359 (1.4)	2471 (97.9)		
Arrival timing				
am	16057 (98.5)	245 (1.5)	0.39	0.52
pm	11598 (98.6)	166 (1.4)		

N.B.: * χ^2 ; **Mann-Whitney test

Table 3. Relationship between boarding time and patients' characters, arrival method, eligibility, outcome, triage level, fate, and comorbidities

Variable	Boarding Time				Test	p-value
	< 2 hours	2- < 6 hours	6-12 hours	> 12 hours		
Age	34.28 ± 10.78	34.07 ± 20.54	34.06 ± 20.79	33.85 ± 22.58	3**	0.47
Gender						
Female	7123 (51.2)	4288 (30.8)	2132 (15.3)	376 (2.7)	9.01*	0.02
Male	7126 (50.4)	4523 (32)	2068 (14.6)	430 (3.0)		
Nationality						
Saudi	11244 (50.6)	6894 (31.2)	3327 (15.1)	630 (2.9)	2.26*	0.51
Non - Saudi	3005 (50.3)	1917 (32.1)	873 (14.6)	176 (2.9)		
Triage level						
Priority 1 - Resuscitation	166 (50.8)	99 (30.3)	48 (14.7)	14 (4.3)	9.2*	0.66
Priority 2 - Emergent	2042 (20.9)	1216 (30.3)	628 (15.7)	122 (3.0)		
Priority 3 - Urgent	4043 (50.3)	2570 (32)	1200 (14.9)	226 (2.8)		
Priority 4 - Less Urgent	7554 (51.0)	4642 (31.4)	2185 (14.8)	424 (2.9)		
Priority 5 - Non-Urgent	444 (50.1)	284 (32)	139 (15.7)	20 (2.3)		
Arrival method						
Ambulance	120 (52.4)	62 (27.1)	39 (17.0)	8 (3.5)	2.46*	0.48
Other	14129 (50.8)	8749 (31.4)	4161 (14.9)	798 (2.9)		
Eligibility						
No	1299 (51.5)	761 (30.2)	379 (15)	84 (3.3)	2.64*	0.3
Yes	12950 (50.7)	8050 (31.5)	3821 (15)	722 (2.8)		
Arrival timing						
am	8176 (50.2)	5179 (31.8)	2470 (15.2)	477 (2.9)	5.95	0.11
pm	6073 (51.6)	3632 (30.9)	1730 (14.7)	329 (2.8)		

N.B.: * χ^2 ; **Mann-Whitney test



N.B.: * χ^2 ; **Mann-Whitney test

Figure 2. Relationship between patients' death and boarding time.

Table 4. Binary logistic regression analysis regarding the risk factors for death

Variable	Depression			p-value
	Beta	Wald	Exp. (B)	
Boarding time	1.86	839.82	6.46	< 0.001
Eligibility	0.36	5.08	1.43	0.02

DISCUSSION

The current research aimed to assess the association between boarding time in the ED and increased patient mortality at the King Abdul-Aziz University Hospital. Patients who died had a significantly longer boarding time than those who survived, indicating a highly significant link between death and boarding time. This study found that 46.2% of patients who died had a boarding time of more than 12 hours compared to 2.2% for those who did not die, with a statistically significant difference. Our results are similar to previous studies, which found lower mortality among patients transferred out of the ED in less than 6 hours^[22,23,24,25].

About half of the participants (50.8%) had a boarding duration of less than two hours, while just 2.9% had a boarding time of more than 12 hours. Patients who were sicker and required more resource-intensive inpatient care had the greatest ED waiting time, according to a prior study. These excessive wait times are linked to a much longer length of stay in the hospital^[19]. Another study found that patients who were in the ED for more than 4 hours before being transferred to an inpatient bed had a higher risk of dying in the hospital^[17].

Singer et al. discovered a similar link between patient fatality and boarding time. This study discovered that hospitalized patients who spent more than four hours in the emergency room were at a higher risk of serious health problems and mortality^[26]. Intas et al. discovered that ED boarding for longer than 6 hours was linked to a higher risk of death in the hospital^[27]. Patients presenting to the emergency department during shifts with longer waiting times were linked to a higher risk of death and hospital admission in the near term, according to a study conducted in the United States^[7].

Our study showed that patients with orthopedic or neurological disorders and injuries had a significantly higher percentage of boarding time of more than 12 hours (5.2%). This is because these patients require

more time for stabilization and imaging. These findings are consistent with a study by Nori et al. that showed that the mean stay time in the ED for boarded and non-boarded orthopedic patients was 1.07 hours and 10.18 hours, respectively^[28].

In the present study, 52.8% of the patients were non-emergency cases, and 99.2% were walk-in patients to the ED, which is considered the primary cause of ED overcrowding. This matter can affect health caregivers, care recipients, health supplies, and eventually, the country's economy^[29]. Primary health care, staff clinics, family physicians, urgent care centers, and even home healthcare services can help in most circumstances. The current study found that 52.8% of the patients were non-emergency cases, and 99.2% of them were walk-in patients to the ED, which is the main cause of ED overcrowding. This issue impacts healthcare providers, patients, medical supplies, and the country's economy^[30]. However, alternatives such as primary health care, staff clinics, family physicians, urgent care centers, and home healthcare services can help in most situations. In Saudi Arabia, citizens have complete access to KAUH without any restrictions to receive free medical treatment for people of all ages, and medical services are available to everyone equally. The Ministry of Health serves primary health care through a network of regional health centers^[31]. For the past 20 years, the government has funded new and innovative programs to ensure health facilities are available to all citizens at all care levels including primary, secondary, and tertiary centers like KAUH ED, which accept referrals from various areas.

Furthermore, non-Saudi tourists accounted for 21% of all ED visits. According to current Saudi legislation, any health insurance must cover all non-Saudis (excluding sponsored individuals) before performing their legal papers. Health insurance in Saudi Arabia requires the prior submission of papers, which includes all nationalities (excluding individual sponsorships)^[30, 32]. However, a previous study found that the sickest hospital patients are "frequent flyers." Their admission

rates are the highest, their death rates are the highest, and their resource consumption is disproportionately high^[33].

Various strategies can be used to alleviate ED overpopulation. Bedside registration and effective utilization of hospitalists are examples of internal departmental advances. Additional solutions include provider triage and provider scheduling optimization. Adding beds to the emergency room does not always solve the problem of boarding and overcrowding. Externally, smoothing elective (schedulable) admissions is likely the single most important strategy for increasing capacity, reducing boarding, maintaining nurse-to-patient ratios, and improving ICU access. Early morning discharges from inpatient units also substantially decrease ED boarding^[34, 35]. Increased weekend discharges and improved weekend services will increase capacity and reduce boarding. When an institution's bed capacity is exceeded, a full-capacity strategy, in which inpatient units go over the census rather than boarding patients in the ED, is safer, preferable to patients, and reduces the duration of stay.

LIMITATIONS

Considering the limitations of the study design, a cause-and-effect relationship is difficult to establish. A retrospective analysis of the data also carries the risk of missing data or inaccurate entries.

CONCLUSION

This study discovered a significant association between patient death and boarding time, with patients who died having a significantly longer boarding time than those who survived. About 46% of patients who died had an average boarding time of more than 12 hours, compared to 2.2 percent of those who did not. The death rate was significantly higher among non-eligible patients and male patients, and those with injuries or orthopedic diseases had a significantly higher percentage of those who had more than 12 hours of boarding time. Long-term boarding and non-eligibility for management were both risk factors for death among the patients investigated. The impact of protracted boarding times on patient outcomes is highlighted in this paper, highlighting the necessity for immediate steps and attempts to decrease ED boarding.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare. All co-authors have seen and agreed with the contents of the manuscript. There are no financial interests to disclose. We certify that the submission is an original work and is not under review at any other publication.

DISCLOSURE

The authors did not receive any form of commercial support, either in the form of compensation or financial assistance, for this case report. The authors have no financial interest in any of the products, devices, or drugs mentioned in this article.

ETHICAL APPROVAL

This is a retrospective study for which the consent requirement was waived. The study was conducted in accordance with the Declaration of Helsinki, and the Ethics Committee of King Abdulaziz University approved the protocol (Ethical Approval Reference No 281-18).

REFERENCES CITED

- [1] Trzeciak S, Rivers EP. Emergency department overcrowding in the United States: an emerging threat to patient safety and public health. *Emerg Med J.* 2003;20(5):402-405.
- [2] Wang Z, Xiong X, Wang S, et al. Causes of emergency department overcrowding and blockage of access to critical services in Beijing: a 2-year study. *J Emerg Med.* 2018;54(5):665-673.
- [3] McCarthy ML, Ding R, Zeger SL, et al. A randomized controlled trial of the effect of service delivery information on patient satisfaction in an emergency department fast track. *Acad Emerg Med.* 2011;18(7):674-685.
- [4] Pines JM, Iyer S, Disbot M, et al. The effect of emergency department crowding on patient satisfaction for admitted patients. *Acad Emerg Med.* 2008;15(9):825-831.
- [5] Bernstein SL, Aronsky D, Duseja R, et al. The effect of emergency department crowding on clinically oriented outcomes. *Acad Emerg Med.* 2009;16(1):1-10.
- [6] Chalfin DB, Trzeciak S, Likourezos A, et al. Impact of delayed transfer of critically ill patients from the emergency department to the intensive care unit. *Crit Care Med.* 2007;35(6):1477-1483.
- [7] Guttmann A, Schull MJ, Vermeulen MJ, et al. Association between waiting times and short term mortality and hospital admission after departure from emergency department: population based cohort study from Ontario,

- Canada. *BMJ*. 2011;342:d2983.
- [8] Hollander JE, Pines JM. The emergency department crowding paradox: the longer you stay, the less care you get. *Ann Emerg Med*. 2007;50(5):497-499.
- [9] Pines JM, Hollander JE. Emergency department crowding is associated with poor care for patients with severe pain. *Ann Emerg Med*. 2008;51(1):1-5.
- [10] Schneider S, Winograd S. Emergency department crowding. *Emergency Medicine Reports*. 2009;30(3):13-23.
- [11] Shakhathreh H, Karim A, Al-Durgam M, et al. Use and misuse of accident and emergency services at Queen Alia Military Hospital. *JR Med Serv*. 2003;10(1):44-47.
- [12] Siddiqui S, Ogbeide DO. Utilization of emergency services in a community hospital. *Saudi Med J*. 2002;23(1):69-72.
- [13] Horwitz LI, Green J, Bradley EH. US emergency department performance on wait time and length of visit. *Ann Emerg Med*. 2010;55(2):133-141.
- [14] Schull MJ, Vermeulen M, Slaughter G, et al. Emergency department crowding and thrombolysis delays in acute myocardial infarction. *Ann Emerg Med*. 2004;44(6):577-585.
- [15] Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345(19):1368-1377.
- [16] Gordon JA, Billings J, Asplin BR, et al. Safety net research in emergency medicine proceedings of the academic emergency medicine consensus conference on "The Unraveling Safety Net". *Acad Emerg Med*. 2001;8(11):1024-1029.
- [17] Boulain T, Malet A, Maitre O. Association between long boarding time in the emergency department and hospital mortality: a single-center propensity score-based analysis. *Intern Emerg Med*. 2020;15(3):479-489.
- [18] McHugh M, VanDyke K, McClelland M, et al. Improving patient flow and reducing emergency department crowding: a guide for hospitals. 2012. <https://www.ahrq.gov/sites/default/files/publications/files/ptflowguide.pdf>
- [19] Salehi L, Phalpher P, Valani R, et al. Emergency department boarding: a descriptive analysis and measurement of impact on outcomes. *CJEM*. 2018;20(6):929-937.
- [20] Diercks DB, Roe MT, Chen AY, et al. Prolonged emergency department stays of non-ST-segment-elevation myocardial infarction patients are associated with worse adherence to the American College of Cardiology/American Heart Association guidelines for management and increased adverse events. *Ann Emerg Med*. 2007;50(5):489-496.
- [21] Al-Khathaami AM, Abulaban AA, Mohamed GE, et al. The impact of 'admit no bed' and long boarding times in the emergency department on stroke outcome. *Saudi Med J*. 2014;35(9): 993-998.
- [22] Intas G, Stergiannis P, Chalari E, et al. The impact of ED boarding time, severity of illness, and discharge destination on outcomes of critically ill ED patients. *Adv Emerg Nurs J*. 2012; 1;34(2):164-169.
- [23] Khan BA, Shakeel N, Siddiqui EU, et al. Impact of delay in admission on the outcome of critically ill patients presenting to the emergency department of a tertiary care hospital from low income country. *J Pak Med Assoc*. 2016;66(5):509-516.
- [24] Teklie H, Engida H, Melaku B, et al. Factors contributing to delay intensive care unit admission of critically ill patients from the adult emergency department in Tikur Anbessa Specialized Hospital. *BMC Emerg Med*. 2021; 26;21(1):123-132.
- [25] Hung SC, Kung CT, Hung CW, et al. Determining delayed admission to intensive care unit for mechanically ventilated patients in the emergency department. *Crit Care*. 2014;23;18(4):485-494.
- [26] Singer AJ, Thode HC Jr, Viccellio P, et al. The association between length of emergency department boarding and mortality. *Acad Emerg Med*. 2011;18(12):1324-1329.
- [27] Intas G, Stergiannis P, Chalari E, et al. The impact of ED boarding time, severity of illness, and discharge destination on outcomes of critically ill ED patients. *Adv Emerg Nurs J*. 2012; 1;34(2):164-169.
- [28] Nouri Y, Gholipour C, Aghazadeh J, et al. Evaluation of the risk factors associated with emergency department boarding: a retrospective cross-sectional study. *Chin J Traumatol*. 2020;23(6):346-350. doi: 10.1016/j.cjtee.2020.09.002/
- [29] Dawoud SO, Ahmad AM, Alsharqi OZ, et al. Utilization of the emergency department and predicting factors associated with its use at the Saudi Ministry of Health General Hospitals. *Glob J Health Sci*. 2015;17;8(1):90-106.
- [30] Almalki M, Fitzgerald G, Clark M. Health care system in Saudi Arabia: an overview. *East Mediterr Health J*. 2011;17(10):784-793.
- [31] Saudi Arabia International Travel Information. U.S. Department of State Bureau of Consular Affairs. 2021. <https://travel.state.gov/content/travel/en/international-travel/International-Travel-Country-Information-Pages/SaudiArabia.html>
- [32] Huang M, van der Borgh C, Leithaus M, et al. Patients' perceptions of frequent hospital admissions: a qualitative interview study with older people above 65 years of age. *BMC Geriatr*. 2020; 7;20(1):332-344.
- [33] Sinclair D. Emergency department overcrowding - implications for paediatric emergency medicine. *Paediatr Child Health*. 2007;12(6):491-494.
- [34] Salway RJ, Valenzuela R, Shoenberger JM, et al. Emergency department (ed) overcrowding: evidence-based answers to frequently asked questions. *RMCLC*. 2017;28(2):213-219.
- [35] McKenna P, Heslin SM, Viccellio P, et al. Emergency department and hospital crowding: causes, consequences, and cures. *Clin Exp Emerg Med*. 2019;6(3):189-195.